



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



Publication number: **0 247 983 B1**

(12)

## EUROPEAN PATENT SPECIFICATION

(45) Date of publication of patent specification: 07.01.93 (51) Int. Cl.<sup>5</sup>: **A61K 31/44, A61K 9/30, A61K 9/54**

(21) Application number: **87850127.9**

(22) Date of filing: **16.04.87**

Divisional application 92107179.1 filed on  
16/04/87.

(54) **New pharmaceutical preparation for oral use.**

(30) Priority: **30.04.86 GB 8610572**

(43) Date of publication of application:  
**02.12.87 Bulletin 87/49**

(45) Publication of the grant of the patent:  
**07.01.93 Bulletin 93/01**

(84) Designated Contracting States:  
**AT BE CH DE ES FR GB GR IT LI LU NL SE**

(56) References cited:  
**EP-A- 0 080 602 EP-A- 0 124 495**  
**DE-A- 3 233 764 FR-A- 2 272 639**  
**GB-A- 760 403 GB-A- 1 190 387**

**PATENT ABSTRACTS OF JAPAN, vol. 8, no.**  
**106 (C-223)[1543], 18 May 1984\***

**SCAND. J. GASTROENTEROLOGY, vol.20,**  
**supplement 108, 1985\***

**BROCHURE on HPMC TC-5 (Shinetsu Chemi-**  
**cal)\***

**UP TO DATE PHARMACEUTICAL TECHNOL-**  
**OGY, series no. 1: Coating of Drugs, 1969\***

(73) Proprietor: **Aktiebolaget Hässle**  
**Kärragatan 5**  
**S-431 83 Mölndal(SE)**

(72) Inventor: **Lövgren, Kurt Ingmar**  
**Violinvägen 2D**  
**S-435 00 Mölnlycke(SE)**  
Inventor: **Pilbrant, Ake Gunnar**  
**Snödroppevägen 6**  
**S-434 00 Kungsbacka(SE)**  
Inventor: **Yasumura, Mitsuru**  
**37, Matsuzono 5-chome**  
**Nishinomiya Hyogo 662(JP)**  
Inventor: **Morigaki, Satoshi**  
**275-84, Nakanoma Yachlyo-cho**  
**Taka-gun Hyogo 677-01(JP)**  
Inventor: **Oda, Minoru**  
**304-24, Ohaza-Yuya**  
**Nakatsu-shi Ohita 871(JP)**  
Inventor: **Ohishi, Naohiro**  
**1342-1, Ohaza-Tarumi Shinyoshitomimura**  
**Chikujo-gun Fukuoka 871-09(JP)**

(74) Representative: **Linderöth, Margareta et al**  
**AB Astra Patent Department**  
**S-151 85 Södertälje(SE)**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

EP 0 247 983 B1

Int. J. OF PHARMACEUTICS, vol. 31, 1986; pp.  
131-136\*

ACTA CHEMICA SCANDINAVICA, vol. 43,  
1989, no. 6; pp. 536-611 (reprint)\*

---

**Description**

The present invention is related to a new stable pharmaceutical preparation containing omeprazole for oral use, and to a method for the manufacture of such a preparation.

From e.g. EP-A1-0 005 129 omeprazole, 5-methoxy-2-(((4-methoxy-3,5-dimethyl-2-pyridinyl)methyl)-sulfinyl)-1H-benzimidazole, a potent inhibitor of gastric acid secretion is known. Omeprazole shows a powerful inhibitory action against secretion of gastric juice (Lancet, Nov 27, 1982, p. 1223-1224) and can be used for the treatment of gastric and duodenal ulcers. Omeprazole is however susceptible to degradation/transformation in acid reacting and neutral media. The half-life of omeprazole in water solutions at pH-values less than four is shorter than ten minutes. Also at neutral pH-values the degradation reaction proceeds rapidly, e.g. at pH=7 the half-life of omeprazole is about 14 hours, while at higher pH-values the stability in solution is much better (Pilbrant and Cederberg, Scand. J. Gastroenterology 1985; 20 (suppl. 108) p. 113-120). The stability profile is similar in solid phase. The degradation of omeprazole is catalyzed by acidic reacting compounds and is stabilized in mixtures with alkaline reacting compounds. The stability of omeprazole is also affected by moisture and organic solvents.

From what is said about the stability properties of omeprazole, it is obvious that an oral dosage form of omeprazole must be protected from contact with the acid reacting gastric juice in order to reach the small intestine without degradation.

In human pharmacological studies it was found that the rate of release of omeprazole from a pharmaceutical dosage form can influence the total extent of absorption of omeprazole to the general circulation (Pilbrant and Cederberg, Scand. J. Gastroenterology 1985; 20 (suppl. 108) p. 113-120). A fully bioavailable dosage form of omeprazole must release the active drug rapidly in the proximal part of the gastrointestinal canal.

In order to obtain a pharmaceutical dosage form of omeprazole which prevents omeprazole from contact with acidic gastric juice, the cores must be enteric coated. Ordinary enteric coatings, however, are made of acidic compounds. If covered with such a conventional enteric coating, omeprazole rapidly decomposes by direct or indirect contact with it, with the result that the preparations become badly discolored and lose in omeprazole content with the passage of time.

In order to enhance the storage stability the cores which contain omeprazole must also contain alkaline reacting constituents. When such an alkaline core is enteric coated with an amount of a conventional enteric coating polymer such as, for example, cellulose acetate phthalate, that permits the dissolution of the coating and the active drug contained in the cores in the proximal part of the small intestine, it also will allow some diffusion of water of gastric juice through the enteric coating into the cores, during the time the dosage form resides in the stomach before it is emptied into the small intestine. The diffused water of gastric juice will dissolve parts of the core in the close proximity of the enteric coating layer and there form an alkaline solution inside the coated dosage form. The alkaline solution will interfere with the enteric coating and eventually dissolve it.

An enteric coated dosage form of omeprazole was reported by Pilbrant and Cederberg, in the above cited Scand. J. Gastroenterology 1985; 20 (suppl. 108) p. 113-120. The publication describes a conventional enteric coated dosage form and states that it has an acceptable storage stability - for clinical studies. It was later found that the stability of this dosage form was insufficient during long-term storage required for a marketed pharmaceutical dosage form.

If a conventional formulation of omeprazole is made, the stability is not satisfactory, particularly in resistance to humidity, and special moisture-proof packing has been adopted to minimize the troubles. However, this provides no satisfactory solution to the problems in today's drug distribution system, and also leads to increased costs. Under the circumstances, there has been a demand for the development of new enteric preparations of omeprazole with better stability.

In DE-A1-3046 559 a way to coat a dosage form is described. First the dosage form is coated with a water insoluble layer containing microcrystalline cellulose and then with a second enteric coating with the aim to achieve a dosage form which releases the active drug in the colon. This method of preparation will not give the desired release of omeprazole in the small intestine.

US-A-2 540 979 describes an enteric coated oral dosage form, where the enteric coating is combined with a second and/or first coating of a water insoluble "wax" layer. This method of preparation is not applicable on cores containing omeprazole since direct contact between substances such as cellulose acetate phthalate (CAP) and omeprazole causes degradation and discolouration of omeprazole.

DE-B2-23 36 218 describes a method to produce a dialysis membrane consisting of a mixture of one or more conventional enteric coating polymers and one or more insoluble cellulose derivatives. Such a membrane will not give a proper protection of omeprazole in gastric juice.

DE-A1-1 204 363 describes a three-layer coating procedure. The first layer is soluble in gastric but is insoluble in intestinal juice. The second is water soluble regardless of pH and the third layer is an enteric coating. This preparation as well as the preparation described in DE-A1-1 617 615 result in a dosage form which is not dissolved in gastric juice and which only dissolves slowly in intestinal juice. Such preparations cannot be used for omeprazole, where a rapid release of the drug in the small intestine is needed.

DE-A1 12 04 363 describes coating with three layers to achieve release of a drug in the ileum, an aim which is outside the scope of the present invention.

GB-A-1 485 676 describes a way to obtain a preparation, which effervesces in the small intestine, by enteric coating a core containing the active drug and an effervescing system such as a combination of carbonate and/or bicarbonate salt and a pharmaceutically acceptable acid. The formulation cannot be adopted for a pharmaceutical dosage form containing omeprazole, as the presence of an acid in contact with omeprazole in the cores would give a result that omeprazole was degraded.

WO 85/03436 describes a pharmaceutical preparation, wherein cores containing active drugs mixed with for instance buffering components such as sodium dihydrogenphosphate with the aim of maintaining a constant pH and a constant rate of diffusion, are coated with a first coating which controls the diffusion. This formulation cannot be adopted for omeprazole where a rapid release in the small intestine is wanted. Direct application of an enteric coating onto the cores would also adversely influence the storage stability of such dosage forms containing omeprazole.

EP-A-124 495 describes enteric coated granules without subcoating or powder that are filled into hard gelatine capsules or a solution that is filled into soft capsules.

The object of the present invention is to provide an enteric coated dosage form of omeprazole, which is stable to discolouration and which is resistant to dissolution in acid media and which dissolves rapidly in neutral to alkaline media and which has a good stability during long-term storage. The new dosage form is characterized in the following way. Core material in the form of small beads or tablets containing omeprazole together with an alkaline reacting compound, or an alkaline salt of omeprazole optionally together with an alkaline reacting compound, and on said core material one or more inert reacting subcoating layers comprising tablet excipients which are soluble or rapidly disintegrating in water, or polymeric, water soluble, filmforming compounds, optionally containing pH-buffering, alkaline compounds between the alkaline reacting core and an outer layer, which is an enteric coating. This/these inner layer/layers separates/separate the alkaline core material from the outer layer, which is an enteric coating. The final, enteric coated dosage form is treated in a suitable way to reduce the water content to a very low level in order to obtain a good stability of the dosage form during long-term storage.

#### Detailed description of the invention

##### Cores

Omeprazole is mixed with inert, preferably water soluble, conventional pharmaceutical constituents to obtain the preferred concentration of omeprazole in the final mixture and with an alkaline reacting, otherwise inert, pharmaceutically acceptable substance (or substances), which creates a "micro-pH" around each omeprazole particle of not less than pH=7, preferably not less than pH=8, when water is adsorbed to the particles of the mixture or when water is added in small amounts to the mixture. Such substances can be chosen among, but are not restricted to substances such as the sodium, potassium, calcium, magnesium and aluminium salts of phosphoric acid, carbonic acid, citric acid or other suitable weak inorganic or organic acids; substances normally used in antacid preparations such as aluminium, calcium and magnesium hydroxides; magnesium oxide or composite substances, such as  $A_1_2O_3 \cdot 6MgO \cdot CO_2 \cdot 12H_2O$ ,  $(Mg_6A_1_2(OH)_{16}CO_3 \cdot 4H_2O)$ ,  $MgO \cdot A_1_2O_3 \cdot 2SiO_2 \cdot nH_2O$  or similar compounds; organic pH-buffering substances such as trihydroxymethylaminomethane or other similar, pharmaceutically acceptable pH-buffering substances. The stabilizing, high pH-value in the powder mixture can also be achieved by using an alkaline reacting salt of omeprazole such as the sodium, potassium, magnesium, calcium etc. salts of omeprazole, which are described in e.g. EP-A2-124 495, either alone or in combination with a conventional buffering substance as previously described.

The powder mixture is then formulated into small beads i.e. pellets, or tablets, by conventional pharmaceutical procedures. The pellets or tablets are used as cores for further processing. The pellets or tablets are used as cores for further processing.

##### Separating layer

The omeprazole containing alkaline reacting cores must be separated from the enteric coating polymer(s) containing free carboxyl groups, which otherwise causes degradation/discholoration of omeprazole during the coating process or during storage. The subcoating layer, in the following defined as the separating layer, also serves as a pH-buffering zone in which hydrogen ions diffusing from the outside in towards the alkaline core can react with hydroxyl ions diffusing from the alkaline core towards the surface of the coated particles. The pH-buffering properties of the separating layer can be further strengthened by introducing in the layer substances chosen from a group of compounds usually used in antacid formulations such as, for instance, magnesium oxide, hydroxide or carbonate, aluminium or calcium hydroxide, carbonate or silicate; composite aluminium/magnesium compounds such as, for instance  $\text{Al}_2\text{O}_3 \cdot 6\text{MgO} \cdot \text{CO}_2 \cdot 12\text{H}_2\text{O}$ ,  $(\text{Mg}_6\text{Al}_2(\text{OH})_{16}\text{CO}_3 \cdot 4\text{H}_2\text{O})$ ,  $\text{MgO} \cdot \text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2 \cdot n\text{H}_2\text{O}$  or similar compounds; or other pharmaceutically acceptable pH-buffering compounds such as, for instance the sodium, potassium, calcium, magnesium and aluminium salts of phosphoric, citric or other suitable, weak, inorganic or organic acids.

The separating layer consists of one or more water soluble inert layer, optionally containing pH-buffering compounds.

The separating layer(s) can be applied to the cores - pellets or tablets - by conventional coating procedures in a suitable coating pan or in a fluidized bed apparatus using water and/or conventional organic solvents for the coating solution. The material for the separating layer is chosen among the pharmaceutically acceptable, water soluble, inert compounds or polymers used for film-coating applications such as, for instance sugar, polyethylene glycol, polyvinylpyrrolidone, polyvinyl alcohol, hydroxypropyl cellulose, methylcellulose, hydroxymethyl cellulose, hydroxypropyl methylcellulose or polyvinyl acetate diethylaminoacetate. The thickness of the separating layer is not less than 2  $\mu\text{m}$ , for small spherical pellets preferably not less than 4  $\mu\text{m}$ , for tablets preferably not less than 10  $\mu\text{m}$ .

In the case of tablets another method to apply the coating can be performed by the drycoating technique. First a tablet containing omeprazole is compressed as described above. Around this tablet a layer is compressed using a suitable tableting machine. The outer, separating layer, consists of pharmaceutically acceptable, in water soluble or in water rapidly disintegrating tablet excipients. The separating layer has a thickness of not less than 1 mm. Ordinary plasticizers, colorants, pigments, titanium dioxide, talc and other additives may also be included into the separating layer.

### 30 Enteric coating layer

The enteric coating layer is applied on to the sub-coated cores by conventional coating techniques such as, for instance, pan coating or fluidized bed coating using solutions of polymers in water and/or suitable organic solvents or by using latex suspensions of said polymers. As enteric coating polymers can be used, for example, cellulose acetate phthalate, hydroxypropyl methylcellulose phthalate, polyvinyl acetate phthalate, carboxymethylcellulose, co-polymerized methacrylic acid/methacrylic acid methyl esters such as, for instance, compounds known under the trade name Eudragit® L 12,5 or Eudragit® L 100 (Röhm Pharma), or similar compounds used to obtain enteric coatings. The enteric coating can also be applied using water-based polymer dispersions, e.g. Aquateric® (FMC Corporation), Eudragit® L100-55 (Röhm Pharma), Coating CE 5142 (BASF). The enteric coating layer can optionally contain a pharmaceutically acceptable plasticizer such as, for instance, cetanol, triacetin, citric acid esters such as, for instance, those known under the trade name Citroflex® (Pfizer), phthalic acid esters, dibutyl succinate or similar plasticizers. The amount of plasticizer is usually optimized for each enteric coating polymer(s) and is usually in the range of 1-20 % of the enteric coating polymer(s). Dispersants such as talc, colorants and pigments may also be included into the enteric coating layer.

Thus, the special preparation according to the invention consists of cores containing omeprazole mixed with an alkaline reacting compound or cores containing an alkaline salt of omeprazole optionally mixed with an alkaline reacting compound. The alkaline reacting core material and/or alkaline salt of the active ingredient, omeprazole, enhance the stability of omeprazole. The cores suspended in water forms a solution or a suspension which has a pH, which is higher than that of a solution in which the polymer used for enteric coating is just soluble. The cores are coated with an inert reacting water soluble or in water rapidly disintegrating coating, optionally containing a pH-buffering substance, which separates the alkaline cores from the enteric coating. Without this separating layer the time during which the dosage form maintains the resistance towards gastric juice would be too short and/or the storage stability of the dosage form would be unacceptably short. The sub-coated dosage form is finally coated with an enteric coating rendering the dosage form insoluble in acid media, but rapidly disintegrating/dissolving in neutral to alkaline media such as, for instance the liquids present in the proximal part of the small intestine, the site where dissolution is wanted.

Final dosage form

The final dosage form is either an enteric coated tablet or enteric coated pellets, pellets dispensed in hard gelatin capsules or sachets, or enteric coated pellets formulated into tablets. It is essential for the long term stability during storage that the water content of the final dosage form containing omeprazole (enteric coated tablets, or pellets) is kept low, preferably not more than 1.5 % by weight. As a consequence the final package containing hard gelatin capsules filled with enteric coated pellets preferably also contain a desiccant, which reduces the water content of the gelatin shell to a level where the water content of the enteric coated pellets filled in the capsules does not exceed 1.5 % by weight.

Process

A process for the manufacture of the oral dosage form represents a further aspect of the invention. After the forming of the cores the cores are first coated with the separating layer(s) and then with the enteric coating layer. The coating is carried out as described above.

The preparation according to the invention is especially advantageous in reducing gastric acid secretion and/or providing a gastrointestinal cytoprotective effect. It is administered one to several times a day. The typical daily dose of the active substance varies and will depend on various factors such as the individual requirements of the patients, the mode of administration and disease. In general the daily dose will be in the range of 1-400 mg of omeprazole.

The invention is described in detail in the following examples:

EXAMPLESExample 1

The effect of different magnesium compounds was evaluated in the form of enteric coated tablets. Tablet cores were first made by known techniques according to the formulations listed in Table 1, followed by application of separating layers and enteric coating layers as shown in Table 2.

Table 1

Formulations for the tablet cores (mg)							
Formulations No.	1	2	3	4	5	6	7
Omeprazol	15.0	15.0	15.0	15.0	15.0	15.0	15.0
Lactose	134.0	119.0	119.0	119.0	118.8	118.5	119.0
Hydroxypropyl cellulose (low substitution	5.0	5.0	5.0	5.0	5.0	5.0	5.0
Hydroxypropyl cellulose	1.0	1.0	1.0	1.0	1.0	1.0	1.0
Talc	5.0	5.0	5.0	5.0	5.0	5.0	5.0
Na <sub>2</sub> HPO <sub>4</sub>	-	15.0	-	-	0.2	-	-
Na lauryl sulfate	-	-	-	-	-	0.5	-
MgO	-	-	15.0	-	-	-	-
Mg(OH) <sub>2</sub>	-	-	-	15.0	15.0	15.0	-
Synthetic hydrotalcite [Al <sub>2</sub> O <sub>3</sub> • 6MgO • CO <sub>2</sub> • 12H <sub>2</sub> O]	-	-	-	-	-	-	15.0
Total	160.0	160.0	160.0	160.0	160.0	160.0	160.0

Table 2 Formulations for coatings (mg)

Formulation No.	I	II	III	IV
5 Separating layer (inner):				
Hydroxypropyl cellulose	-	2.0	2.0	2.0
Magnesium hydroxide	-	-	0.3	-
10 Synthetic hydrotalcite	-	-	-	0.3
Separating layer (outer):				
Hydroxypropyl cellulose	-	2.0	2.0	2.0
Enteric coating layer:				
15 Hydroxypropyl methylcellulose				
phthalate	7.0	7.0	7.0	7.0
Cetyl alcohol	0.5	0.5	0.5	0.5

20 The tablets thus obtained were stored in open form under so called accelerated conditions, that is 40° C, and 75 % relative humidity, and the changes in appearance with the passage of time were observed. Storage for six months under these conditions corresponds to storage at normal temperature for three years. This means that high stability sufficient for practical use may be assured if a drug remains intact for  
 25 about one week under the mentioned conditions. The result is summarized in Table 3. As may be seen from the table, a remarkable stabilizing effect is achieved when a magnesium compound is contained in the inner separating layer.

Table 3

Stabilizing Effect (Appearance of Preparations)								
Coating Layer		Core material						
		1	2	3	4	5	6	7
I	At the start	C	A	A	A	A	A	A
	60° C; after 7 days	E	D	C	C	C	C	D
	40° C; 75 % RH; after 7 days	F	E	B	B	B	B	E
II	At the start	A	A	A	A	A	A	A
	60° C; after 7 days	E	B	A	A	A	A	C
	40° C; 75 % RH; after 7 days	E	D	A	A	A	A	D
III	At the start	A	A	A	A	A	A	A
	60° C; after 15 days	B	A	A	A	A	A	A
	40° C; after 30 days	A	A	A	A	A	A	A
	40° C; 75 % RH; after 15 days	B	A	A	A	A	A	A
IV	At the start	A	A	A	A	A	A	A
	60° C; after 15 days	B	A	A	A	A	A	A
	40° C; after 30 days	A	A	A	A	A	A	A
	40° C; 75 % RH; after 15 days	B	A	A	A	A	A	A
A: white, B: brownish white, C: faint brown, D: light brown, E: brown, F: deep brown.								

55 All the samples evaluated as A (white) in the above table showed no discoloration even on split surfaces. The samples evaluated as B (brownish white) showed little change in appearance, but some discoloration was observed on split surfaces.

Table 4 shows the result of a stability test on the omeprazole preparation according to Example 1 (Formulation No 4-IV). The formulation was stored in a closed glass bottle at room temperature for the

# EP 0 247 983 B1

indicated period of time. This clearly demonstrates that preparations with unusually high stability were obtained.

Table 4

Stability of enteric coated omeprazole preparations (Tablets of Formulation No. 4-IV)		
Storage Period	Appearance	Omeprazole Content (%)
At the start of test	White	100.0
1 year at room temperature	White	99.9
2 years at room temperature	White	100.0

## Example 2

### Uncoated pellets

I	Mannitol powder	16 150 g
	Lactose anhydrous	800 g
	Hydroxypropyl cellulose	600 g
	Microcrystalline cellulose	400 g
II	Omeprazole	2 000 g
	Sodium lauryl sulphate	50 g
	Disodium hydrogen phosphate	80 g
	Distilled water	4 400 g

The dry ingredients (I) were premixed in a mixer. Addition of a granulation liquid (II) containing suspended omeprazole was made and the mass was wet-mixed to a proper consistency. The wet mass was pressed through an extruder and spheronized to pellets. The pellets were dried and classified into suitable particle size ranges.

### Subcoated pellets

III	Uncoated omeprazole pellets	6 000 g
	Hydroxypropyl methylcellulose	240 g
	Distilled water	4 800 g

The polymer solution (III) was sprayed on the uncoated pellets in a fluidized bed apparatus. The spray guns were placed above the fluidized bed.

### Enteric-coated pellets



		Subcoated pellets	500 g
5	IV	{ Hydroxypropyl methylcellulose	
		{ phthalate	57 g
		{ Cetyl alcohol	3 g
		{ Acetone	540 g
10		{ Ethanol	231 g

The polymer solution (IV) was sprayed on the subcoated pellets in a fluidized bed apparatus with spray guns placed above the bed. After drying to a water content of 0.5 % the enteric coated pellets were classified and filled into hard gelatin capsules in an amount of 225 mg, corresponding to 20 mg of omeprazole. 30 capsules were packed in tight containers together with a desiccant.

### Example 3

This example illustrates that a variety of polymers can be used for subcoating, e.g. hydroxypropyl methylcellulose, hydroxypropyl cellulose, polyvinylpyrrolidone, polyethylene glycol, polyvinyl alcohols.

### Uncoated pellets

25		{ Mannitol powder	1 620 g
		{ Lactose anhydrous	80 g
30	I	{ Hydroxypropyl cellulose	60 g
		{ Microcrystalline cellulose	40 g
		{ Omeprazole	200 g
35		{ Sodium lauryl sulphate	1.0 g
	II	{ Disodium hydrogen phosphate	9.3 g
		{ Distilled water	515 g
40			

The uncoated pellets were prepared as described in Example 2.

### Subcoated pellets

45		{ Uncoated omeprazole pellets	500 g
	III	{ Polyvinylpyrrolidone	20 g
50		{ Ethanol	400 g

The subcoated pellets were prepared as described in Example 2.

### Enteric-coated pellets

5		{	Subcoated pellets	500 g
			Hydroxypropyl methyl-	
			cellulose phthalate	45 g
10	IV	{	Cetyl alcohol	5 g
			Acetone	219 g
			Ethanol	680 g

The enteric-coated pellets were prepared as described in Example 2.

Example 4

Uncoated pellets

20		{	Mannitol powder	1 610 g
			Lactose anhydrous	80 g
25	I	{	Hydroxypropyl cellulose	60 g
			Microcrystalline cellulose	40 g
30		{	Omeprazole	200 g
			Pluronic® F68	10 g
	II	{	Disodium hydrogen phosphate	24 g
35			Distilled water	450 g

The uncoated pellets were prepared as described in Example 2.

Subcoated pellets

40			Uncoated pellets	500 g
45	III	{	Polyvinylpyrrolidone	30 g
			Ethanol	400 g

The subcoated pellets were prepared as described in Example 2.

Enteric coated pellets

		Subcoated pellets	500 g
5		Hydroxypropyl methyl-	
		cellulose phthalate	45 g
	IV	Cetyl alcohol	5 g
		Methylene chloride	371 g
10		Ethanol	680 g

The enteric coated pellets were prepared as described in Example 2.

15

#### Example 5

This example illustrates that a variety of polymers can be used as enteric coating material e.g. cellulose acetate phthalate, poly-(vinyl acetate/vinyl alcohol phthalate), hydroxypropyl methylcellulose phthalate, poly-  
 20 (methacrylic acid/methacrylic acid methyl esters), poly-(acrylic acid/methacrylic acid methyl esters). The polymers can be applied with/without plasticizer, e.g. polyethylene glycols, triacetin, dimethyl polysiloxan, Citroflex®, cetyl alcohol, stearyl alcohol, diethyl phthalate.

Enteric-coated pellets can also be manufactured from water-based polymer dispersions, e.g. Aquateric® (FMC Corporation), Eudragit®L 100-55, Coating CE 5142 (BASF).

25

#### Uncoated pellets

30		Lactose powder	277 g
		Lactose anhydrous	118 g
	I	Hydroxypropyl cellulose	25 g
35		Colloidal silica	25 g
		Omeprazole	50 g
		Sodium lauryl sulphate	5 g
40	II	Disodium hydrogen phosphate	2 g
		Sodium dihydrogen phosphate	0.1 g
45		Distilled water	170 g

The uncoated pellets were prepared as described above.

50

#### Subcoated pellets

The uncoated pellets were subcoated as described in Example 2.

#### Enteric coated pellets

55

5	III	{	Subcoated pellets	500 g
			Eudragit® L 100	45 g
			Stearyl alcohol	4.5 g
			Ethanol	1 320 g

10

The enteric coated pellets were prepared as described above.

#### Example 6

15 Formulations with the sodium salt of omeprazole.

#### Uncoated pellets

20		{	Omeprazole sodium salt	339 g
			Mannitol powder	2 422 g
			Lactose anhydrous	120 g
25	I	{	Hydroxypropyl cellulose	90 g
			Microcrystalline cellulose	60 g
30	II	{	Sodium lauryl sulphate	7 g
			Distilled water	650 g

35 The preparation was made as described in Example 2 with the exception that the omeprazole sodium salt was added together with the other ingredients in mixture I.

#### Subcoated pellets

40			Uncoated pellets	500 g
	III	{	Hydroxypropyl methylcellulose	20 g
45			Aluminium hydroxide/magnesium carbonate	4 g
			Distilled water	400 g
50	IV	{	Pellets subcoated with III	500 g
			Hydroxypropyl methylcellulose	20 g
			Distilled water	400 g

55

The two subcoat layers, III and IV, were applied to the uncoated pellets in a fluidized bed apparatus in consecutive order as previously described.

Enteric coated pellets

		Subcoated pellets	500 g
5	V	Hydroxypropyl methylcellulose	
		phthalate	57 g
		Cetyl alcohol	3 g
10		Acetone	540 g
		Ethanol	231 g

15 The preparation of enteric coated pellets was performed as described in Example 2.

Examples 7 and 8

Formulations with the magnesium salt of omeprazole.

20

Uncoated pellets

			<u>Example No</u>	
25			<u>7</u>	<u>8</u>
	I	Omeprazole magnesium salt	222 g	222 g
30		Mannitol powder	1 673 g	1 473 g
		Microcrystalline cellulose	100 g	100 g
		Magnesium hydroxide	-	200 g
35	II	Sodium lauryl sulphate	5 g	5 g
		Distilled water	500 g	375 g

40

The preparation was made as described in Example 2 with the exception that the omeprazole magnesium salt was added together with the other ingredients in mixture I.

Subcoated pellets

45

			<u>Example</u>	
50			<u>7 and 8</u>	
	III	Uncoated pellets	500 g	
55		Hydroxypropyl methyl-		
		cellulose	20 g	
		Distilled water	400 g	

The pellets were prepared as described in Example 2.

#### Enteric coated pellets

5

#### Examples

#### 7 and 8

10

Subcoated pellets 500 g

15

IV {	Hydroxypropyl methyl-	
	cellulose phthalate	57 g
	Cetyl alcohol	3 g
	Acetone	540 g
	Ethanol	231 g

20

The enteric coated pellets were prepared as described in Example 2.

#### Examples 9 and 10

Manufacture of tablets.

#### Tablet cores

30

#### Examples No

#### 9      10

35

I {	Omeprazole	400 g	-
	Omeprazole sodium salt, corre-		
	sponding to omeprazole 400 g	-	426 g
	Lactose, anhydrous	1 420 g	1 409 g
	Polyvinylpyrrolidone,		
crosslinked	100 g	100 g	
	Sodium carbonate, anhydrous	15 g	-

45

II {	Methyl cellulose	12 g	12 g
	Distilled water	200 g	200 g

50

Magnesium stearate	30 g	30 g
--------------------	------	------

55

The powder mixture I was carefully homogenized and granulated by the solution II. The wet mass was dried in a fluidized bed dryer using an inlet air temperature of +50°C for 30 minutes. The dried mixture

## EP 0 247 983 B1

was then forced through a sieve with an aperture of 0.5 mm. After mixing with magnesium stearate the granulate was tabletted on a tableting machine using 6 mm punches. The tablet weight was 100 mg.

### Subcoating

The tablets containing omeprazole were subcoated with approximately 10 % by weight of hydroxypropyl methylcellulose from a water solution using a perforated coating pan apparatus.

The tablets containing omeprazole sodium salt were subcoated using the dry coating technique. A tablet granulate containing

Lactose anhydrous	4 000 g
Polyvinylpyrrolidone, (PVP)	180 g
Ethanol 95 %	420 g
Magnesium stearate	42 g

was prepared in the following way. The lactose was granulated with a solution of PVP in ethanol and dried. After drying magnesium stearate was admixed.

The granulate mass was dry coated around the tablet cores of example 9 using a Manesty Dry Cota® tableting machine. The tablet weight of the dry coated tablets was 475 mg. Each tablet contained 20 mg of omeprazole.

### Enteric coating

The subcoated tablets obtained above were enteric coated using the same coating solution:

Hydroxypropyl methylcellulose phthalate	1 500 g
Cetyl alcohol	105 g
Methylene chloride	15 000 g
Isopropanol	15 000 g
Distilled water	3 150 g

The coating was applied in a perforated coating pan apparatus. An approximate amount of one kg of coating solution was applied for each kg of tablets.

## COMPARATIVE EXAMPLES

### Examples I, II and III

These examples illustrate that the buffer salt used effects the enteric-coated omeprazole pellets properties when the sub-coating layer is absent. A high amount of buffer salt is needed in order to obtain a long shelf life for the product. At the same time this type of pellets shows inferior acid resistance properties. C.f. also the Example 4 above.

### Uncoated pellets

		Examples No		
		I	II	III
5				
	I			
	{ Mannitol powder	1 610 g	1 610 g	1 610 g
	{ Lactose anhydrous	80 g	80 g	80 g
10	{ Hydroxypropyl			
	{ cellulose	60 g	60 g	60 g
	{ Microcrystalline			
15	{ cellulose	40 g	40 g	40 g
	II			
	{ Omeprazole	200 g	200 g	200 g
	{ Pluronic® F68	10 g	10 g	10 g
20	{ Disodium hydrogen			
	{ phosphate	2 g	8 g	24 g
	{ Distilled water	450 g	450 g	450 g
25				

The uncoated pellets were prepared as described in Example 2 above.

30 Enteric coated pellets

		Uncoated pellets	500 g
35	III	{ Hydroxypropyl methylcellulose	
		{ phthalate	45 g
		{ Cetyl alcohol	5 g
		{ Methylene chloride	371 g
40		{ Ethanol	680 g

45 The coated pellets were prepared as described in Example 2 above.

Example IV

This formulation is the same as in Example 6 above, but no subcoating layer was used.

50 Uncoated pellets

55



5	I	Omeprazole sodium salt	339 g
		Mannitol powder	2 422 g
		Lactose anhydrous	120 g
		Hydroxypropyl cellulose	90 g
		Microcrystalline cellulose	60 g
10	II	Sodium lauryl sulphate	7 g
		Distilled water	650 g

15

The preparation was made as described in Example 6.

Enteric-coated pellets

20

	III	Uncoated pellets	500 g
		Hydroxypropyl methylcellulose	
25		phthalate	57 g
		Cetyl alcohol	3 g
		Acetone	540 g
30		Ethanol	231 g

35

The enteric coated pellets were prepared as described in Example 2.

Example V

This formulation is the same as in Example 8 above, but no subcoating layer was used.

Uncoated pellets

45	I	Omeprazole magnesium salt	222 g
		Mannitol powder	1 473 g
		Microcrystalline cellulose	100 g
		Magnesium hydroxide	200 g
50	II	Sodium lauryl sulphate	5 g
		Distilled water	375 g

55

The preparation was made as described in Example 8.

Enteric coated pellets

EP 0 247 983 B1

	Uncoated pellets	500 g
	Hydroxypropyl methylcellulose	
5	phthalate	57 g
	Cetyl alcohol	3 g
	Acetone	540 g
10	Ethanol	231 g

The pellets were prepared as described in Example 2 above.

15 Properties of the enteric coated pellets

For the preparations according to Examples 2-8 and comparative Examples I-V above one or both of the following studies have been performed.

20 Acid resistance

The resistance of the formulations was studied in the following way: The formulations were added to gastric fluid USP (without enzyme), 37 °C (paddle) 100 r/min. After 2 hours the actual amount of omeprazole remaining intact in the formulations was determined.

25 Rate of dissolution in buffer solution

In order to establish the rate of dissolution in the small intestine, the formulations were added to a buffer solution. Buffer solution 37 °C, USP dissolution apparatus No 2 (paddle), 100 r/min. After 10 or 30 minutes the amount of omeprazole dissolved was determined. The results are presented in the following Table 5.

35

40

45

50

55

Table 5

Example No	Omeprazole content mg/g	Acid resistance, amount intact omeprazole (%)		% dissolved omeprazole at different pH:s and after 10 or 30 min	
		after 2 hours	%	pH	min
2	89.2	95	100	6.8	10
3	90	96	91	6.0	10
4	88	89	*)		
5	82	93	70	7.5	30
6	81.3	87	93	6.8	10
7	91	95	**)		
8	89	98	**)		
I	93	97	*)		
II	92	94	*)		
III	94	58	*)		
IV	86.5	4	*)		
V	91	93	**)		

\*) The stability of the formulation was studied during storage in glass bottles also containing a desiccant device. After one month storage at +50 °C the formulation according to Example 4 was virtually intact with no change in appearance or physicochemical characteristics. Pellets according to Example I and II turned brown due to degradation, while the pellets according to Example III retained to original white colour.

\*\*) The formulations according to Examples 7 and 8 were white and not affected by the coating process. The enteric coated pellets according to Example V, where the enteric coating was applied directly on the cores according to Example 8, was discoloured already during the enteric coating process.

Further comparative test

5 This example demonstrates the effect of the moisture content of the preparations according to the invention on storage stability.

The stability of omeprazole pellets according to the invention was compared with that of omeprazole pellets with higher water content. Omeprazole pellets were prepared according to the invention with a water content of 1 %. Two other portions of the same formulation were conditioned to a water content of 2 % and 5 % respectively. The three formulations, packed in tight containers not containing a desiccant, were stored for one month at +50 °C. After this time the packages were opened and the pellets were assayed for the amount of omeprazole by HPLC. The formulation according to the invention had an omeprazole content of 98.5 % of the initial value. The other two formulations with a water content of 2 and 5 % respectively were virtually totally degraded and had only trace amounts of intact omeprazole.

15 DISCUSSION

From the results given in Table 5 it can be seen that formulations containing omeprazole with acceptable acid resistance can be prepared by using a conventional enteric coating technique (see for instance Examples I, II and V). However, it is also obvious that the storage stability of the formulations according to Examples I, II and V is not acceptable, since a discolouration, showing a degradation of omeprazole, occurs during short storage at an elevated storage temperature (Examples I and II) or already during the enteric coating process (Example V).

If the amount of alkaline substances in the cores is increased to a level where omeprazole has an acceptable storage stability (Example III) or if an alkaline reacting salt of omeprazole is used in the preparation of the cores (Example IV), then, without the separating layer of the invention, the resistance to dissolution in acid media becomes unacceptably low and much or all of the active substance will degrade already in the stomach and thus, it has no effect on the gastric acid secretion.

When the preparation is carried out according to the invention as for instance in Example 4, a good resistance towards gastric juice as well as a good stability during long-term storage is obtained. This is in contrast with the formulations in Examples I, II and III where either an acceptable acid resistance or an acceptable storage stability can be achieved - but not both. The same comparison can be made between the formulations according to Examples 7 and 8 according to the invention and the formulation according to Example V, where the separating layer was omitted. Examples 7 and 8 differ in that a buffering substance, magnesium hydroxide, has been included in the cores of Example 8. This further improves the acid resistance as well as the storage stability of Example 8 in comparison with Example 7.

The further comparative test shows the great importance of a low water content in the preparations.

Thus in order to prepare pharmaceutical formulations of omeprazole for oral use, which exert good stability during long-term storage as well as good stability during the residence in the stomach after administration, the preparation is made in the following way:

- a) Omeprazole together with an alkaline reacting compound or compounds or an alkaline reacting salt of omeprazole optionally mixed with alkaline reacting compound are included in the core material.
- b) The core material is subcoated with one or more inert, in water soluble or in water rapidly disintegrating layers, which separate the alkaline reacting core from the enteric coating. The subcoating layer may optionally contain pH-buffering compounds.
- c) The subcoated cores are coated with an acid insoluble enteric coating, optionally containing plasticizers.

Biopharmaceutical studies

10 The hard gelatin capsules according to Example 2 were administered to 12 healthy, young male volunteers in the following way:

The volunteers came to the laboratory in the morning after having abstained from food since 10 p.m. the night preceding the experimental day. A zero time blood sample was taken. One omeprazole capsule according to Example 2 was administered together with 150 ml of tap water. Further blood samples were taken during the day.

In another experiment the same volunteers were administered 20 mg of omeprazole in the form of a suspension of micronized omeprazole in a sodium bicarbonate water solution. In order to reduce the

degradation of omeprazole in the stomach to a minimum, sodium bicarbonate solution was given to the subjects just before the administration of the omeprazole suspension and at further four times with a 10-minutes interval after the drug intake. The concentration of omeprazole in blood plasma was assayed by high pressure liquid chromatography (Persson, Lagerström and Grundevik. Scand J Gastroenterol 1985, 20, (suppl 108), 71-77. The mean plasma concentrations are given in Table 6.

**Table 6**

The plasma concentrations ( $\mu\text{mol/l}$ ) after 20 mg single oral doses of omeprazole given as hard gelatin capsules according to Example 2 and as a suspension of micronized omeprazole in sodium bicarbonate solution.

Time (min)	Capsules	Suspension
10		0.84
20		0.90
30	0.03	0.84
45		0.64
60	0.22	0.44
90	0.36	0.24
120	0.39	0.13
150	0.29	
180	0.20	0.04
210	0.10	
240	0.05	0.01
300	0.02	0
360	0.01	
420	0	

Although the plasma concentration peak at different times, the two formulations are bioequivalent. The mean relative bioavailability of the capsules in comparison with the suspension was 85 %  $\pm$  23 % (S.D.). The comparison was based on the total area under individual plasma concentration versus time curves.

Thus, by preparing capsules according to the invention it is possible to obtain a preparation with the same bioavailability as a suspension containing the same amount of micronized active compound. It is, however, to be noticed that when the suspension is administered, the patients must also be given sodium bicarbonate solution frequently in order to minimize pre-absorption degradation of omeprazole in the stomach.

#### Claims

Claims for the following Contracting States : BE, CH, DE, FR, GB, IT, LI, LU, UL, SE

1. An oral, pharmaceutical preparation stable to discolouration containing omeprazole as the active ingredient characterized in that it is composed of core material in the form of small beads or tablets containing omeprazole together with an alkaline reacting compound, or an alkaline salt of omeprazole optionally together with an alkaline reacting compound, and on said core material one or more inert reacting subcoating layers comprising tablet excipients which are soluble or rapidly disintegrating in

## EP 0 247 983 B1

water, or polymeric, water soluble, filmforming compounds, optionally containing pH-buffering, alkaline compounds between the alkaline reacting core and an outer layer, which is an enteric coating.

2. A preparation according to claim 1 wherein the subcoating comprises hydroxypropyl methylcellulose, hydroxypropyl cellulose or polyvinylpyrrolidone.
3. A preparation according to claim 1 wherein the subcoating comprises two or more sub-layers and where the inner sublayer contains one or more of magnesium oxide, magnesium hydroxide or composite substance  $\text{Al}_2\text{O}_3 \cdot 6\text{MgO} \cdot \text{CO}_2 \cdot 12\text{H}_2\text{O}$  or  $\text{MgO} \cdot \text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2 \cdot n\text{H}_2\text{O}$ , wherein n is not an integer and less than 2.
4. A preparation according to claim 1 wherein the alkaline core comprises omeprazole and pH-buffering alkaline compound rendering to the micro-environment of omeprazole a pH of 7-12.
5. A preparation according to claim 4 wherein the alkaline compound comprises one or more of magnesium oxide, hydroxide or carbonate, aluminium hydroxide, aluminium, calcium, sodium or potassium carbonate, phosphate or citrate, the composite aluminium/magnesium compounds  $\text{Al}_2\text{O}_3 \cdot 6\text{MgO} \cdot \text{CO}_2 \cdot 12\text{H}_2\text{O}$  or  $\text{MgO} \cdot \text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2 \cdot n\text{H}_2\text{O}$ , where n is not an integer and less than 2.
6. A preparation according to claim 1 wherein the alkaline core comprises an alkaline salt of omeprazole such as the sodium, potassium, magnesium, calcium or ammonium salt.
7. A preparation according to claim 6 wherein the alkaline core comprises an alkaline salt of omeprazole mixed with an alkaline compound.
8. A preparation according to claim 1 wherein the enteric coating comprises hydroxypropyl methylcellulose phthalate, cellulose acetate phthalate, co-polymerized methacrylic acid/methacrylic acid methyl ester or polyvinyl acetate phthalate, optionally containing a plasticizer.
9. A preparation according to claim 1 wherein the water content of the final dosage form containing omeprazole does not exceed 1.5 % by weight.
10. Process for the preparation of an oral pharmaceutical formulation stable to discolouration containing omeprazole according to claim 1 in which cores containing omeprazole mixed with an alkaline reacting compound or compounds or an alkaline salt of omeprazole optionally mixed with an alkaline reacting compound or compounds are coated with one or more inert reacting subcoating layers whereafter the subcoated cores are further coated with an enteric coating.

### Claims for the following Contracting States : AT, ES, GR

1. A process for the preparation of an oral, pharmaceutical formulation stable to discolouration containing omeprazole as the active ingredient characterized in that the omeprazole mixed with an alkaline reacting compound, or an alkaline salt of omeprazole optionally mixed with an alkaline reacting compound, are formed to cores and said cores, which are in the form of small beads or tablets are coated with one or more inert reacting subcoating layers comprising tablet excipients which are soluble or rapidly disintegrating in water, or polymeric water soluble, filmforming compounds, optionally containing pH-buffering, alkaline compounds between the alkaline reacting core and an outer layer, which is an enteric coating layer, whereafter the subcoated cores are further coated with said outer enteric coating layer.
2. A process according to claim 1 wherein the applied subcoating comprises hydroxypropyl methylcellulose, hydroxypropyl cellulose or polyvinylpyrrolidone.
3. A process according to claim 1 wherein the applied subcoating comprises two or more sub-layers and where the inner sub-layer contains one or more of magnesium oxide, magnesium hydroxide or composite substance  $\text{Al}_2\text{O}_3 \cdot 6\text{MgO} \cdot \text{CO}_2 \cdot 12\text{H}_2\text{O}$  or  $\text{MgO} \cdot \text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2 \cdot n\text{H}_2\text{O}$ , wherein n is not an integer and less than 2.

4. A process according to claim 1 wherein omeprazole is mixed with a pH-buffering alkaline compound rendering to the micro-environment of omeprazole a pH of 7-12 to form an alkaline core.
5. A process according to claim 4 wherein the alkaline compound which omeprazole is mixed with comprises one or more of magnesium oxide, hydroxide or carbonate, aluminium hydroxide, aluminium, calcium, sodium or potassium carbonate, phosphate or citrate, the composite aluminium/magnesium compounds  $\text{Al}_2\text{O}_3 \cdot 6\text{MgO} \cdot \text{CO}_2 \cdot 12\text{H}_2\text{O}$  or  $\text{MgO} \cdot \text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2 \cdot n\text{H}_2\text{O}$ , where n is not an integer and less than 2.
- 10 6. A process according to claim 1 wherein the alkaline salt of omeprazole such as the sodium, potassium, magnesium, calcium or ammonium salt is formed and incorporated into the alkaline core.
7. A process according to claim 6 wherein the alkaline salt of omeprazole is mixed with an, alkaline compound, to form the alkaline core.
- 15 8. A process according to claim 1 wherein the enteric coating which comprises hydroxypropyl methylcellulose phthalate, cellulose acetate phthalate, co-polymerized methacrylic acid/methacrylic acid methyl ester or polyvinyl acetate phthalate, optionally containing a plasticizer is applied.
- 20 9. A process according to claim 1 wherein a dosage form containing omeprazole is prepared in which the water content does not exceed 1.5 % by weight.

#### Patentansprüche

Patentansprüche für folgende Vertragsstaaten : BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

- 25 1. Orales, pharmazeutisches Präparat, das gegen Verfärbung stabil ist und Omeprazol als das aktive Ingrediens enthält, dadurch gekennzeichnet, daß es aus Kernmaterial in Form kleiner Kügelchen oder Tabletten, welches Omeprazol zusammen mit einer alkalisch reagierenden Verbindung, oder einem alkalischen Omeprazol-Salz, gegebenenfalls zusammen mit einer alkalisch reagierenden Verbindung,
  - 30 enthält, und aus einer oder mehreren inert reagierenden Basisüberzugsschichten auf diesem Kernmaterial, umfassend in Wasser lösliche oder darin rasch zerfallende Tablettenträgermittel, oder polymere, wasserlösliche, filmbildende Verbindungen, die gegebenenfalls pH-puffernde, alkalische Verbindungen zwischen dem alkalisch reagierenden Kern und einer äußeren Schicht, die ein enterischer Überzug ist, enthalten, zusammengesetzt ist.
- 35 2. Präparat nach Anspruch 1, worin der Basisüberzug Hydroxypropylmethylcellulose, Hydroxypropylcellulose oder Polyvinylpyrrolidon umfaßt.
3. Präparat nach Anspruch 1, worin der Basisüberzug zwei oder mehrere Unterschichten aufweist und
  - 40 wobei die innere Unterschicht eines oder mehrere von Magnesiumoxid, Magnesiumhydroxid oder der zusammengesetzten Substanz  $\text{Al}_2\text{O}_3 \cdot 6\text{MgO} \cdot \text{CO}_2 \cdot 12\text{H}_2\text{O}$  oder  $\text{MgO} \cdot \text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2 \cdot n\text{H}_2\text{O}$ , worin n keine ganze Zahl und weniger als 2 ist, umfaßt.
- 45 4. Präparat nach Anspruch 1, worin der basische Kern Omeprazol und eine pH-puffernde alkalische Verbindung umfaßt, die der Mikroumgebung von Omeprazol einen pH-Wert von 7-12 verleiht.
5. Präparat nach Anspruch 4, worin die alkalische Verbindung eines oder mehrere von Magnesiumoxid, -hydroxid oder -carbonat, Aluminiumhydroxid, Aluminium-, Calcium, Natrium- oder Kaliumcarbonat, -phosphat oder -citrat, der zusammengesetzten Aluminium/Magnesiumverbindungen
  - 50  $\text{Al}_2\text{O}_3 \cdot 6\text{MgO} \cdot \text{CO}_2 \cdot 12\text{H}_2\text{O}$  oder  $\text{MgO} \cdot \text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2 \cdot n\text{H}_2\text{O}$ , worin n keine ganze Zahl und weniger als 2 ist, umfaßt.
6. Präparat nach Anspruch 1, worin der alkalische Kern ein alkalisches Omeprazol-Salz, wie das Natrium-, Kalium-, Magnesium-, Calcium- oder Ammoniumsalz, umfaßt.
- 55 7. Präparat nach Anspruch 6, worin der alkalische Kern ein alkalisches Omeprazol-Salz in Mischung mit einer alkalischen Verbindung umfaßt.

8. Präparat nach Anspruch 1, worin der enterische Überzug Hydrosypropylmethylcellulosephthalat, Celluloseacetatphthalat, co-polymerisierten Methacrylsäure/Methacrylsäuremethylester oder Polyvinylacetatphthalat umfaßt und gegebenenfalls einen Weichmacher enthält.
- 5 9. Präparat nach Anspruch 1, worin der Wassergehalt der endgültigen, Omeprazol enthaltenden Dosisform 1,5 Gew.-% nicht übersteigt.
10. Verfahren zur Herstellung einer oralen pharmazeutischen Formulierung, die gegen Verfärbung stabil ist und Omeprazol enthält, nach Anspruch 1, bei welchem Kerne, die Omeprazol gemischt mit einer alkalisch reagierenden Verbindung oder Verbindungen oder ein alkalisches Salz von Omeprazol, gegebenenfalls gemischt mit einer alkalisch reagierenden Verbindung oder Verbindungen enthalten, mit einer oder mehreren inert reagierenden Basisüberzugsschichten überzogen werden, wonach die mit dem Basisüberzug überzogenen Kerne weiters mit einem enterischen Überzug beschichtet werden.
- 15 **Patentansprüche für folgende Vertragsstaaten : AT, ES, GR**
1. Verfahren zur Herstellung einer oralen, pharmazeutischen Formulierung, die gegen Verfärbung stabil ist und Omeprazol als das aktive Ingrediens enthält, dadurch gekennzeichnet, daß das mit einer alkalisch reagierenden Verbindung gemischte Omeprazol oder ein alkalisches Omeprazol-Salz, gegebenenfalls in Mischung mit einer alkalisch reagierenden Verbindung, zu Kernen geformt wird und diese Kerne, die in Form kleiner Kügelchen oder Tabletten vorliegen, mit einer oder mehreren inert reagierenden Basisüberzugsschichten umfassend in Wasser lösliche oder darin rasch zerfallende Tablettenträgermittel, oder polymere wasserlösliche filmbildende Verbindungen, die gegebenenfalls pH-puffernde, alkalische Verbindungen zwischen dem alkalisch reagierenden Kern und einer äußeren Schicht, die eine enterische Überzugsschicht ist, enthalten, überzogen werden, wonach die mit dem Basisüberzug überzogenen Kerne weiters mit der äußeren enterischen Überzugsschicht beschichtet werden.
- 20 2. Verfahren nach Anspruch 1, worin der aufgebrauchte Basisüberzug Hydroxypropylmethylcellulose, Hydroxypropylcellulose oder Polyvinylpyrrolidon umfaßt.
3. Verfahren nach Anspruch 1, worin der aufgebrauchte Basisüberzug zwei oder mehrere Unterschichten umfaßt, und wobei die innere Unterschicht eines oder mehrere von Magnesiumoxid, Magnesiumhydroxid oder der zusammengesetzten Substanz  $\text{Al}_2\text{O}_3 \cdot 6\text{MgO} \cdot \text{CO}_2 \cdot 12\text{H}_2\text{O}$  oder  $\text{MgO} \cdot \text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2 \cdot n\text{H}_2\text{O}$ , worin n keine ganze Zahl und weniger als 2 ist, umfaßt.
- 35 4. Verfahren nach Anspruch 1, worin Omeprazol mit einer pH-puffernden alkalischen Verbindung gemischt wird, die der Mikroumgebung von Omeprazol einen pH-Wert von 7-12 verleiht, um einen basischen Kern zu bilden.
- 40 5. Verfahren nach Anspruch 4, worin die alkalische Verbindung, mit der Omeprazol gemischt wird, eines oder mehrere von Magnesiumoxid, -hydroxid oder -carbonat, Aluminiumhydroxid, Aluminium-, Calcium-, Natrium- oder Kaliumcarbonat, -phosphat oder -citrat, den zusammengesetzten Aluminium/Magnesiumverbindungen  $\text{Al}_2\text{O}_3 \cdot 6\text{MgO} \cdot \text{CO}_2 \cdot 12\text{H}_2\text{O}$  oder  $\text{MgO} \cdot \text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2 \cdot n\text{H}_2\text{O}$ , worin n keine ganze Zahl und weniger als 2 ist, umfaßt.
- 45 6. Verfahren nach Anspruch 1, worin das alkalische Omeprazol-Salz, wie das Natrium-, Kalium-, Magnesium-, Calcium- oder Ammoniumsalz, gebildet und in den alkalischen Kern inkorporiert wird.
7. Verfahren nach Anspruch 6, worin das alkalsiche Salz von Omeprazol mit einer alkalischen Verbindung zur Bildung des alkalischen Kerns gemischt wird.
- 50 8. Verfahren nach Anspruch 1, worin der enterische Überzug der Hydroxypropylmethylcellulosephthalat, Celluloseacetatphthalat, co-polymerisierten Methacrylsäure/Methacrylsäuremethylester oder Polyvinylacetatphthalat umfaßt und gegebenenfalls einen Weichmacher enthält, aufgetragen wird.
- 55 9. Verfahren nach Anspruch 1, worin eine Omeprazol enthaltende Dosisform hergestellt wird, in welcher der Wassergehalt 1,5 Gew.-% nicht übersteigt.



**Revendications**

**Revendications pour les Etats contractants suivants : BE, CH, DE, FR, GB, IT, LI, LU, NL, SE**

1. Préparation pharmaceutique orale stable à la décoloration contenant de l'omeprazole comme ingrédient actif, caractérisée en ce qu'elle est composée d'une matière formant noyau qui se présente sous l'aspect de petites perles ou de comprimés contenant de l'omeprazole conjointement avec un composé à réaction alcaline, ou un sel alcalin d'omeprazole éventuellement conjointement avec un composé à réaction alcaline, et, sur cette matière formant noyau, d'une ou plusieurs sous-couches de revêtement à réaction inerte comprenant des excipients de comprimés qui sont solubles ou se désintègrent rapidement dans l'eau, ou des composés polymères filmogènes solubles dans l'eau, contenant éventuellement des composés alcalins tamponnant le pH entre le noyau à réaction alcaline et une couche externe qui est une couche kératinisée.
2. Préparation selon la revendication 1, dans laquelle la sous-couche de revêtement comprend de l'hydroxypropyl méthylcellulose, de l'hydroxypropyl cellulose ou de la polyvinylpyrrolidone.
3. Préparation selon la revendication 1, dans laquelle la sous-couche de revêtement comprend deux ou plusieurs sous-couches et la sous-couche interne contient une ou plusieurs substances parmi l'oxyde de magnésium, l'hydroxyde de magnésium et une substance composite de formule  $Al_2O_3 \cdot 6MgO \cdot CO_2 \cdot 12H_2O$  ou  $MgO \cdot Al_2O_3 \cdot 2SiO_2 \cdot nH_2O$ , dans laquelle n est un nombre entier qui est inférieur à 2.
4. Préparation selon la revendication 1, dans laquelle le noyau alcalin comprend de l'omeprazole et un composé alcalin tamponnant le pH instaurant un pH de 7-12 dans le micro-environnement de l'omeprazole.
5. Préparation selon la revendication 4, dans laquelle le composé alcalin comprend un ou plusieurs des composés suivants : oxyde, hydroxyde ou carbonate de magnésium, hydroxyde d'aluminium, carbonate, phosphate ou citrate d'aluminium, de calcium, de sodium ou de potassium, des composites d'aluminium et de magnésium de formule  $Al_2O_3 \cdot 6MgO \cdot CO_2 \cdot 12H_2O$  ou  $MgO \cdot Al_2O_3 \cdot 2SiO_2 \cdot nH_2O$  où n n'est pas un nombre entier et est inférieur à 2.
6. Préparation selon la revendication 1, dans laquelle le noyau alcalin comprend un sel alcalin d'omeprazole tel que le sel de sodium, de potassium, de magnésium, de calcium ou d'ammonium.
7. Préparation selon la revendication 6, dans laquelle le noyau alcalin contient un sel alcalin d'omeprazole mélangé à un composant alcalin.
8. Préparation selon la revendication 1, dans laquelle la couche kératinisée comprend du phtalate d'hydroxypropyl méthylcellulose, de l'acétate phtalate de cellulose, du méthyl ester d'acide méthacrylique/acide méthacrylique copolymérisés ou du poly(acétate phtalate de vinyle), contenant éventuellement un plastifiant.
9. Préparation selon la revendication 1, dans laquelle la teneur en eau de la formule de dosage finale contenant l'omeprazole ne dépasse pas 1,5 % en poids.
10. Procédé de préparation d'une formulation pharmaceutique orale stable à la décoloration contenant de l'omeprazole selon la revendication 1, dans lequel les noyaux contenant de l'omeprazole mélangé à un ou des composés à réaction alcaline ou un sel alcalin d'omeprazole éventuellement mélangé à un ou des composés à réaction alcaline sont revêtus d'une ou plusieurs sous-couches de revêtement à réaction inerte, après quoi les noyaux revêtus de ces sous-couches sont encore revêtus d'une couche kératinisée.

**Revendications pour les Etats contractants suivants : AT, ES, GR**

1. Procédé de préparation d'une formulation pharmaceutique orale stable à la décoloration contenant de l'omeprazole comme ingrédient actif, caractérisé en ce que l'omeprazole mélangé à un composé à réaction alcaline ou un sel alcalin d'omeprazole éventuellement mélangé à un composé à réaction

alcaline, est conformé en noyaux et ces noyaux, qui se présentent sous l'aspect de petites perles ou de comprimés, sont revêtus d'une ou plusieurs sous-couches de revêtement à réaction inerte comprenant des excipients de comprimés qui sont solubles ou se désintègrent rapidement dans l'eau, ou des composés polymères filmogènes solubles dans l'eau, contenant éventuellement des composés alcalins tamponnant le pH entre le noyau à réaction alcaline et une couche externe qui est une couche kératinisée, après quoi les noyaux revêtus des sous-couches sont encore revêtus de la couche kératinisée précitée.

2. Procédé selon la revendication 1, dans lequel la sous-couche de revêtement appliquée comprend de l'hydroxypropyl méthylcellulose, de l'hydroxypropyl cellulose ou de la polyvinylpyrrolidone.
3. Procédé selon la revendication 1, dans lequel la sous-couche de revêtement appliquée comprend deux ou plusieurs sous-couches et la sous-couche interne contient une ou plusieurs substances parmi l'oxyde de magnésium, l'hydroxyde de magnésium et une substance composite de formule  $\text{Al}_2\text{O}_3 \cdot 6\text{MgO} \cdot \text{CO}_2 \cdot 12\text{H}_2\text{O}$  ou  $\text{MgO} \cdot \text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2 \cdot n\text{H}_2\text{O}$ , dans laquelle n est un nombre entier qui est inférieur à 2.
4. Procédé selon la revendication 1, dans lequel on mélange l'omeprazole à un composé alcalin tamponnant le pH instaurant un pH de 7-12 pour former un noyau alcalin.
5. Procédé selon la revendication 4, dans lequel le composé alcalin auquel l'omeprazole est mélangé comprend un ou plusieurs des composés suivants : oxyde, hydroxyde ou carbonate de magnésium, hydroxyde d'aluminium, carbonate, phosphate ou citrate d'aluminium, de calcium, de sodium ou de potassium, des composites d'aluminium et de magnésium de formule  $\text{Al}_2\text{O}_3 \cdot 6\text{MgO} \cdot \text{CO}_2 \cdot 12\text{H}_2\text{O}$  ou  $\text{MgO} \cdot \text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2 \cdot n\text{H}_2\text{O}$  où n n'est pas un nombre entier et est inférieur à 2.
6. Procédé selon la revendication 1, dans lequel on forme et on incorpore au noyau alcalin le sel alcalin d'omeprazole tel que le sel de sodium, de potassium, de magnésium, de calcium ou d'ammonium.
7. Procédé selon la revendication 6, dans lequel on mélange le sel alcalin d'omeprazole à un composant alcalin pour former le noyau alcalin.
8. Procédé selon la revendication 1, dans lequel on applique la couche kératinisée qui comprend du phtalate d'hydroxypropyl méthylcellulose, de l'acétate phtalate de cellulose, du méthyl ester d'acide méthacrylique/acide méthacrylique copolymérisé ou du poly(acétate phtalate de vinyle), contenant éventuellement un plastifiant.
9. Procédé selon la revendication 1, dans lequel on prépare une formule de dosage contenant l'omeprazole dans laquelle la teneur en eau ne dépasse pas 1,5 % en poids.